



Results of study on effect of Antischemin preparation on some of the parameters of the protection system and vascular endothelial during the cerebral ischemia model

B Ariuntsetseg¹, B Sarantsetseg², M Ambaga³, D Tserendagva⁴

¹⁻³New Medicine Medical University, Mongolia

⁴Mongolian National University of Medical Sciences, Mongolia

Abstract

Self-regulating mechanism and driving process of the organism has been a subject matter for the science for many years. Then, very simple determination of antioxidant system which is one of the above regulation is that it is unique biological system that keeps alive nature of the cell, origin for the humanity existence and studying changes and activity of the system on Antischemin is one set of the experiment. Research was carried out on ischemia model through methodology of Farkes E et al., 2007 with in the purpose to study effects of Antischemin on superoxidismutase, glutathione, glutathione peroxidase (GSH-Px pmol/ml), and Vascular endothelial growth factor (VEGF) which indicate antioxidant activity. According to the results of the study glutathione (GSH-ng/L) level in cerebral tissue homogenate has decreased by 26-40.1%, superoxide dismutase (SOD) by 25.8-41%, glutathione peroxidase (GSH-px-pmol/ml) 15.4 by 44.5% and by 57.3-60.9% in the plasma respectively on 1st to 21st days after forming cerebral ischemia on trial animals and it proves rapid reduction of endogenous antioxidants in cerebral and cardiac ischemia. And in animals on which 100 mg/kg Antischemin was used, glutathione (GSH-ng/L) has increased by 16.1-27.2% in tissue homogenate, superoxide dismutase (SOD) by 14-20.9%, glutathione peroxidase by up to 14.3-31.1% on 1st to 21st trial days and it ensures condition to provide tissue with high energy substances such as ATP by increasing endogenous antioxidant reserve during oxygen deficiency. Also 16.7-20.95% decrease of VEGF-pg/ml in control group shows unstable metabolism in vascular and tissue causing main factor to damage vessel during cerebral hypoxia and among the group that used Antischemin VEGF-pg/ml has increased by 11.9-17.9% in brain homogenate and by 7.8-10.5% in comparative group, Bilobil, relatively and it indicates that substances influence during new vessel formation even not so strong.

Keywords: antioxidant system, superoxide dismutase, glutathione, glutathione peroxidase, endothelial growth factor

Introduction

Antioxidant system protects against the effects of peroxidation process on fat compounds of tissue membrane, damage and degradation of membrane, tissue disfunction [1] and Antischemin which has positive effects on this system consists of roots of *Astragalus membranaceus*, *Scutellaria baicalensis* Georgi and leaves of *Gingko bilob* that are planted in Mongolia. Antioxidant activity of this medicine has been studied on ischemia model and main causes for vascular endothelial damage, further for unstable metabolism of vital organs including brain and heart during the ischemia include imbalance of electron and proton flows on "9 stepped cycle of proton and electron conductance with in 14 trillion tissues of human body" and oxygen deficiency [2, 3, 4, 6].

It is important to produce medicine which provides preventive and treatment/curing activity and is capable of activating antioxidant system which is unique biological system for the cell/tissue to keep its alive nature, to be used separately and in combination with other medicines on the above disorder at the integrated system level of dependence between many factors.

Purpose

To study Antischemin effects on superoxidismutase, glutathione, glutathione peroxidase (GSH-Px pmol/ml),

GABA / γ -aminobutyric acid/ and Vascular endothelial growth factor (VEGF) that show antioxidant activity during the cerebral ischemia model formed in rat.

Methodology

Disorder model of cerebral ischemia was executed through methodology by Farkes E et al., 2007 [5]. After making 220-280 gram white rat by using ketamine and attaching to the surgical table tightly entered polyethylene tube into trachea with the help of special laryngoscope, connected with small animal ventilator R407 RWD Life Science and allowed breathing by adjusting average breath 94-104 min with volume of 1.8-2.0 cc per minute. Surgery was carried out under aseptic environment with headlamp. After cutting hair along the neck and sterilizing skin area by 5% iodine solution put out left jugular artery and bound by special suture. Fermentation connecting reaction: Following indicators have been determined by taking blood from heart via vacuum tainer after making the trial animal asleep by injecting ketamine hydrochloride on 1st, 3rd, 7th, 14th and 21st days of experiment after forming cerebral ischemia on trial animal through (Enzyme-linked immunosorbent assay) ELISA device, including, SOD /superoxidismutase/, GSH /glutathione/, GSH-Px /glutathionperoxidase/, Vascular endothelial growth

factor (VEGF) in blood plasma and brain tissue [7].

Results

According to our study, superoxidismutase activity contained in the brain tissue has decreased by 32.9% (7.77 ± 0.10 pg/ml in healthy group, 5.21 ± 0.43 pg/ml, $P < 0.001$ in control group) on the 1st day or acute stage, 28.6% (8.04 ± 0.14 pg/ml in healthy group, 5.74 ± 0.17 pg/ml, $P < 0.001$ in control group) on

the 3rd trial day, 25.8% (7.13 ± 0.18 pg/ml in healthy group, 5.29 ± 0.12 pg/ml, $P < 0.05$ in control group) on the 7th day, 36.5% (7.64 ± 0.13 pg/ml in healthy group, 4.85 ± 0.17 pg/ml, $P < 0.001$ in control group) on the 14th day, 41% (6.89 ± 0.28 pg/ml in healthy group, 4.06 ± 0.12 pg/ml, $P < 0.001$ in control group) on the 21st day after forming cerebral ischemia on the white rat compared to the animals in the healthy group.

Table 1: Amount of superoxide dismutase (SOD) contained in the cerebral tissue homogenate and effects of Antischemin on it during the cerebral ischemia disorder model

№	Trial days	SOD contained in the cerebral tissue (pg/ml)			
		Healthy n=6	Control n=6	Antischemin-100 mg/kg n=6	Bilobil-40 mg/kg n=6
1	1 day	7.77 ± 0.12	$5.21 \pm 0.43^*$	$6.53 \pm 0.53^{**}$	5.93 ± 0.36
2	3 days	8.04 ± 0.13	$5.74 \pm 0.17^*$	$7.03 \pm 0.16^{**}$	$7.09 \pm 0.14^{**}$
3	7 days	7.13 ± 0.18	$5.29 \pm 0.12^*$	$6.69 \pm 0.14^{**}$	$6.94 \pm 0.21^{**}$
4	14 days	7.64 ± 0.13	$4.85 \pm 0.17^*$	$5.64 \pm 0.13^{**}$	$6.20 \pm 0.11^{**}$
5	21 days	6.89 ± 0.28	$4.06 \pm 0.12^*$	$6.03 \pm 0.15^{**}$	$5.86 \pm 0.17^{**}$

*- comparison of indicators of control group with healthy group $P \leq 0.05$, $P \leq 0.001$

** - comparison of indicators of treated group with control group $P \leq 0.05$, $P \leq 0.001$

For animals that were treated by Antischemin prepared by condensed infusion of *Astragalus membranaceus*, *Scutellaria baicalensis* Georgi and leaves of *Gingko bilobit* has increased by 20.2% (5.21 ± 0.43 pg/ml in control group, 6.53 ± 0.53 pg/ml, $P < 0.05$ in Antischemin group) on the 1st day, 18.3% (5.74 ± 0.17 pg/ml in control group, 7.03 ± 0.16 pg/ml, $P < 0.05$ in Antischemin group) on the 3rd day, 20.9% (5.29 ± 0.12 pg/ml in control group, 6.69 ± 0.14 pg/ml, $P < 0.05$ in Antischemin group) on the 7th day, 14% (4.85 ± 0.17 pg/ml in control group, 5.64 ± 0.13 pg/ml, $P < 0.05$ in Antischemin group) on the 14th day, 32.6% (4.06 ± 0.123 in control group, 6.03 ± 0.157 pg/ml, $P < 0.001$ in Antischemin group) on the 21st day of the trial compared to the control group.

Superoxide dismutase content was not so different for animals in comparative group on the 1st day compared to the control group and it has increased by 19% (5.74 ± 0.17 pg/ml in control group, 7.09 ± 0.14 pg/ml, $P < 0.05$ in Bilobil group) on the 3rd day, by 23.77% (5.29 ± 0.12 pg/ml in control group, 6.94 ± 0.21 pg/ml, $P < 0.05$ in Bilobil group) on the 7th day, by 21.77% (4.85 ± 0.17 pg/ml in control group, 6.20 ± 0.11 pg/ml, $P < 0.05$ in Bilobil group) on the 14th day, by 30.7% (4.06 ± 0.12 pg/ml in control group, 5.86 ± 0.17 pg/ml, $P < 0.001$ in Bilobil group) on the 21st day of the trial. Generally, it has been observed that Bilobil and Antischemin preparation are similar for increasing activity of SOD.

Table 2: Amount of glutathione (GSHng/L) brain tissue homogenate during the cerebral ischemia model and effects of Antischemin preparation on it

№	Trial days	Glutathione GSH (ng/L)			
		Healthy n=6	Control n=6	Antischemin-100 mg/kg n=6	Bilobil-40 mg/kg n=6
1	1 day	137.19 ± 1.69	121.35 ± 4.98	127.98 ± 1.93	128.35 ± 6.83
2	3 days	129.23 ± 1.3	$95.56 \pm 2.31^*$	$113.94 \pm 2.85^{**}$	$102.43 \pm 1.54^{**}$
3	7 days	119.89 ± 1.27	$77.31 \pm 2.12^*$	81.77 ± 1.09	79.06 ± 1.09
4	14 days	143.79 ± 1.3	$98.15 \pm 1.67^*$	$134.94 \pm 10.74^{**}$	$132.07 \pm 1.93^{**}$
5	21 days	118.59 ± 1.98	$65.11 \pm 0.88^*$	$84.21 \pm 1.30^{**}$	$79.78 \pm 1.01^{**}$

*- comparison of indicators of control group with healthy group $P \leq 0.05$, $P \leq 0.001$

** - comparison of indicators of treated group with control group $P \leq 0.05$, $P \leq 0.001$

On the 1st trial day, in acute period of forming cerebral ischemia on white rat, glutathione content in cerebral tissue homogenate of untreated control group animals was decreased (137.19 ± 1.69 ng/L in healthy group, 121.35 ± 4.98 ng/L in control group) by 11.54%, on the 3rd day by 1.35 times or 26% (129.23 ± 1.3 ng/L in healthy group, 95.56 ± 2.31 ng/L, $P < 0.05$ in control group); on the 7th day by 35.5% (119.89 ± 1.27 ng/L in healthy group, 77.31 ± 2.12 ng/L, $P < 0.001$ in control group); on the 14th trial day by 31.7% (143.79 ± 1.30 ng/L in healthy group, 98.15 ± 1.67 ng/L, $P < 0.001$ in control group); on the 21st day by 40.1% (118.59 ± 1.98 ng/L in healthy group, 65.11 ± 0.88 ng/L, $P < 0.001$ in control group) compared to the indicators of healthy group.

On the other hand, glutathione content in animals that were given 100 mg/kg Antischemin preparation has increased by 5.18% (121.35 ± 4.98 in control group, 127.98 ± 1.93 ng/L in treated group) on the 1st day, by 16.13% (95.56 ± 2.31 in control group, 113.94 ± 2.85 , $P < 0.05$ in treated group) on the 3rd day, by 5.45% (77.31 ± 2.12 in control group, 81.77 ± 1.09 in treated group) on the 7th day, by 27.26% (98.15 ± 1.67 in control group, 134.94 ± 10.74 , $P < 0.001$ in treated group) on the 14th day, by 22.68% (65.11 ± 0.88 in control group, 84.21 ± 1.30 , $P < 0.001$ in treated group) on the 21st trial day compared to control group.

In animals that were treated by Bilobil in comparative group this indicator was increased by 5.45-6.71% on 1st, 3rd, 7th days,

by 25.68% (98.15±1.67 in control group, 132.07±1.93, P<0.001 in Bilobil group) on the 14th day, by 18.38% (65.11±0.88 in control group, 79.78±1.01, P<0.05 in Bilobil

group) on the 21st day compared to untreated control group animals.

Table 3: Amount of glutathione peroxidase (GSH-Px pmol/ml) brain tissue homogenate during the cerebral ischemia model and effects of Antischemin preparation on it

№	Trial days	Glutathione peroxidase GSH-Px pmol/ml			
		Healthy n=6	Control n=6	Antischemin-100 mg/kg n=6	Bilobil-40 mg/kg n=6
1	1 day	9.46±0.2	5.57±0.14*	6.80±0.22**	6.42±0.25**
2	3 days	9.06±0.06	5.46±0.16*	7.06±0.11**	7.23±0.24**
3	7 days	9.14±0.06	7.73±0.16*	9.03±0.19**	9.20±0.22**
4	14 days	9.72±0.14	6.64±0.08*	9.65±0.18**	9.20±0.22**
5	21 days	10.26±0.12	5.69±0.19*	10.16±0.07**	9.76±0.08**

*- comparison of indicators of control group with healthy group P≤ 0.05, P≤ 0.001

**- comparison of indicators of treated group with control group ≤ 0.05, P≤ 0.001

According to the study results, activity of glutathione peroxidase contained in the cerebral tissue homogenate has decreased by 41.1% (P<0.001) in untrated control group animals (5.57±0.14 pmol/ml) compared to healthy group (9.46±0.20 pmol/ml) on the 1st day after forming cerebral ischemia, on the 3rd day by 39.7% (5.46±0.16 pmol/ml in control group, 9.06±0.06 pmol/ml, P<0.001 in healthy group), on the 7th day by 15.4% (7.73±0.16 pmol/ml in control group, 9.14±0.06 pmol/ml, P<0.05 in healthy group), on the 14th day by 31.6% (6.64±0.08 pmol/ml in control group, 9.72±0.14 pmol/ml, P<0.001 in healthy group), on the 21st day by 44.5% (5.69±0.19 pmol/ml in control group, 10.26±0.12 pmol/ml, P<0.001 in healthy group).

And for animals that used Antischemin preparation it has increased by 18.08% (5.57±0.14 pmol/ml in control group, 6.80±0.22 pmol/ml, P<0.05 Antischemin) on the 1st trial day, by 22.66 % (5.46±0.16 pmol/ml control, 7.06±0.11 pmol/ml,

P<0.001 Antischemin) on the 3rd day, by 14.39% (7.73±0.161 control, 9.03±0.19 pmol/ml, P<0.05 Antischemin) on the 7th day, by 31.19% (6.64±0.08 pmol/ml control, 9.65±0.18 pmol/ml, P<0.001 Antischemin) on the 14th day, by 43.99 % (5.69±0.19 pmol/ml control, 10.16±0.07 pmol/ml, P<0.001 Antischemin) on the 21st trial day compared to the control group.

When using comparative preparation Bilobil with dose of 40 mg/kg, glutathione peroxidase level in the cerebral tissue homogenate has increased by 13.23% (control 5.57±0.14 pmol/ml, Bilobil 6.42±0.258 pmol/ml) on the 1st day, by 24.48% (control 5.46±0.16 pmol/ml, Bilobil 7.23±0.24 pmol/ml, P<0.001) on the 3rd day, by 15.97% on the 7th day by 27.8% on the 14th day, by 41.7 % (P<0.001) on the 21st day, respectively compared to the control group and it has proven that both Antischemin preparation and Bilobil has similar function to increase level of glutathione peroxidase.

Table 4: Test results determined by Vascular endothelial growth factor (VEGF)-r Elisa rat kit in rat plasma and cerebral tissue during/on the cerebral ischemia model

Group	3 days pg/ml		7 days pg/ml	
	Blood plasma	Cerebral tissue homogenate (1:10)	Blood plasma	Cerebral tissue homogenate (1:10)
Healthy n=6	93.36±3.15	84.76±0.79	93.36±3.15	84.76±0.79
Control n=9	67.03±0.47	67.04±8.49*	67.39±0.77	70.54±0.97*
Antischemin n=9	79.76±0.76	75.35±0.89	81.66±1.51	80.08±1.30
Bilobil n=9	83.26±1.87	72.78±0.66	80.69±0.94	78.89±1.31

*- comparison of indicators of control group with healthy group P≤ 0.05, P≤ 0.001

**- comparison of indicators of treated group with control group ≤ 0.05, P≤ 0.001

On the 3rd trial day after forming cerebral ischemia on rat Vascular endothelial growth factor (VEGF) level in the cerebral tissue has decreased by 20.9% (in healthy group 84.76±0.79, in control group 67.04±0.49 pg/ml, P<0.001) on the 1st trial day, by 16.7% (in healthy group 84.76±0.79, in control group 70.54±0.97 pg/ml, P<0.001) on the 7th day, for animals treated by Antischemin preparation it has increased by 11% (in control group 67.04±0.49., in Antischemin group 75.35±0.89 pg/ml, P<0.05) on the 3rd trial day, by 11.9% (in control group 70.54±0.97, in Antischemin group 80.08±1.3 pg/ml, P<0.05) on the 7th trial day, respectively and for animals in Bilobil group it has increased by 7.8% (in control group 67.04±0.49 pg/ml, in Bilobil group 72.78±0.66 pg/ml, P<0.05) on the 3rd trial day, by 10.5% (in control group

70.54±0.97 pg/ml, in Bilobil group 78.89±1.31 pg/ml, P<0.05) on the 7th day, respectively.

And Vascular endothelial growth factor (VEGF) level in the serum has decreased 28.2% (in healthy group 93.36±3.15, in control group 67.03±0.47 pg/ml, P<0.001) on th 3rd trial day, by 27.8% (in healthy group 93.36±3.15 in control group 67.39±0.77 pg/ml, P<0.001) on the 7th day, for animals treated by Antischemin preparation it has increased by 15.9% (in control group 67.03±0.47., in Antischemin group 79.76±0.76 pg/ml, P<0.05) on the 3rd trial day, by 17.5% (in control group 67.39±0.77, in Antischemin group 81.66±1.5 pg/ml, P<0.05) on the 7th trial day, respectively and for animals in Bilobil group it has increased by 19.5% (in control group 67.03±0.47 pg/ml, in Bilobil group 83.26±1.87 pg/ml,

$P < 0.05$) on the 3rd trial day, by 16.48% (in control group 67.39 ± 0.77 pg/ml, in Bilobil group 80.69 ± 0.94 pg/ml, $P < 0.05$) on the 7th trial day, respectively.

Discussion

During the subacute and chronic phases formed cerebral ischemia in rat, multiple stepped occurrences of disease are occurred in the cerebral tissue causing oxygen deficiency dependent on ischemia inside the cell, messy peroxidation of fat, substantial formation of oil hydroperoxyl, activation of antioxidant and much consumption of dispensed compounds such as glutathione which keeps peroxidation. Thus, quantity of dispensed compounds are observed as being reduced during the cerebral ischemia model.

Glutathione reserve is reduced as the sickness becomes chronic which in turn fat peroxidation is activated strongly, free radicals are increased with in the cerebral tissue, resulting in reduction of dispensed compounds such as glutathione due to high activation of endogenous antioxidant protection system in the body during the cerebral and heart/cardial ischemia.

Flavonoid and fenolt compounds contained in *Astragalus membranaceus*, *Scutellaria baicalensis* Georgi and leaves of *Gingko bilobdilutes* blood within the micro circulation of blood, decrease peroxidation and positively effect on reserves of internal endogenous antioxidant system during the ischemia. Based on functions of *Astragalus membranaceus*, *Scutellaria baicalensis* Georgi and leaves of *Gingko bilobto* protect nerve cells, [8, 9] eliminate blood stasis, [10, 15] anti-hypoxia activation, [11] effect on cell protection, [12, 16] protection against blood and cerebral loss caused by cerebral edema, [13, 17] potential to be used for inflammation, immune regulation, kidney and heart support, long life expansion as well as many treatments for disease, we have selected the above three plants for our study object and studied antioxidant activation of the preparation.

Conclusion

1. During the 1st to 21st days after forming cerebral ischemia on the trial rat, glutathione (GSH-ng/L) level in cerebral tissue homogenate has decreased by 26-40.1%, superoxide dismutase (SOD) level in the cerebral tissue homogenate by 25.8-41%, glutathione peroxidase (GSH-px-pmol/ml) level in the cerebral tissue homogenate by 15.4-44.5%, by 57.3-60.9% in the plasma in the animals of untreated control group and reserves of endogenous antioxidants rapidly decrease in cerebral and cardiac ischemia. In animals that used 100 mg/kg dose of Antischemin preparation glutathione (GSH-ng/L) level in the tissue homogenate has increased by 16.1-27.2%, superoxide dismutase (SOD) by 14-20.9%, glutathione peroxidase level by 14.3-31.1% during the 1st to 21st days of formation of cerebral ischemia model and it ensures condition to provide cell with high energy compounds such as ATP by increasing reserves of endogenous antioxidants during the oxygen deficiency in the cerebral tissue.
2. 16.7-20.95% decrease of vascular endothelial growth factor (VEGF-pg/ml) in cerebral tissue homogenate of the control group proves creation of many factors for vascular

damage following unstable metabolism within the vascular and tissue environment during the oxygen deficiency in the cerebral tissue and vascular endothelial growth factor (VEGF-pg/ml) in the cerebral tissue homogenate has increased by 11.9-17.9% in the group that used Antischemin preparation, by 7.8-10.5% in comparative or Bilobil group, respectively proving the preparations effect formation of new vein slightly.

Acknowledgement

We would like to acknowledge Research-Innovation Center at "New Medicine Medical University", "Bio-modeling laboratories" and colleagues at Eliza laboratory at Hulj Borjigon hospital, Professor M. Ambaga, Scientific Advisor Professor B.Sarantsetseg. Also special thanks to members of the experimental team, in particular doctoral candidates S.Oyuntsetseg, B.Enebish, S.Munkhbayar and M.Jigidnorov.

References

1. M Ambaga, A.Tumen-Ulzii "The three main lines of membrane-redox potential within the living cells are enclosed circuits of 9-wire proton and electrostatic flux" the value of science and cognitive significance. UB. 2018.
2. M.Ambaga, A.Tumen-Ulzii "The boundary of the three protruding membrane-redox potential in human and animal bodies is a closed circuit of nine protruding protons and electrons." The importance of science and cognition of new theories. UB. 2019.
3. Ambaga M (2017). The membrane-redox potentials three-state line system dependent-full 9 stepped cycle of proton conductance as the universal metabolic formula and the development of all medical thinking during last 3000 years, Asian Journal of Science and technology, vol.08, Issue, 03, pp. 4485-4488, March.
4. Mbagaga M, Tumen-Ulzii A. NCM medicine. Ulaanbaatar. 2018.
5. Scherbak Natalia Sergeevna. Effects and mechanisms of ischemic preconditioning and post-conditioning of the brain. Thesis for the degree of Doctor of Biological Sciences, Sankt Petersburg, 2016.
6. M.Ambaga, A.Tumen-Ulzii NCM medicine. UB, 2017.
7. B.Galindev, Kh.Tserensuren. Laboratory zoology.Ulaanbaatar 2017.
8. Li M, Li H, Fang F, Deng X, Ma S. Astragaloside IV attenuates cognitive impairments induced by transient cerebral ischemia and reperfusion in mice via anti-inflammatory mechanisms. Neurosci Lett. 2017 Feb 3;639:114-119.
9. Luo Y, Qin Z, Hong Z, Zhang X, Ding D, Fu JH, Zhang WD, Chen J. Astragaloside IV protects against ischemic brain injury in a murine model of transient focal ischemia. Neurosci Lett. 2004 Jun 17;363(3):218-23.
10. Dang X, Miao JJ, Chen AQ, Li PChen L, Liang JR, Xie RM, Zhao Y. The antithrombotic effect of RSNK in blood-stasis model rats. J Ethnopharmacol. 2015 Sep 15;173:266-72.
11. Luo Y, Qin Z, Hong Z, Zhang X, Ding D, Fu JH, Zhang WD, Chen J. Astragaloside IV protects against ischemic brain injury in a murine model of transient focal ischemia. Neurosci Lett. 2004 Jun 17;363(3):218-23.

12. Wu X, Zhou W, Wei Q, Chen P, Li Y. Cytoprotective effects of the medicinal herb *Astragalus membranaceus* on lipopolysaccharide-exposed cells. *Mol Med Rep*. 2018 Sep 14
13. Li H, Wang P, Huang F, Jin J, Wu H, Zhang B, Wang Z, Shi H, Wu X. Astragaloside IV protects blood-brain barrier integrity from LPS-induced disruption via activating Nrf2 antioxidant signaling pathway in mice. *Toxicol Appl Pharmacol*. 2018 Feb 1;340:58-66.
14. Li L, Hou X, Xu R, Liu C, Tu M. Research review on the pharmacological effects of astragaloside IV. *Fundam Clin Pharmacol*. 2017 Feb; 31(1):17-36.
15. Kong X, Kong W, Miao G, Zhao S, Chen M, Zheng X, Bai J. Pretreatment with *scutellaria baicalensis* stem-leaf total flavonoid protects against cerebral ischemia/reperfusion injury in hippocampal neurons. *Neural Regen Res*. 2014 Dec 1;9(23):2066-73.
16. Li N, Feng L, Tan Y, Xiang Y, Zhang R, Yang M. Preparation, Characterization, Pharmacokinetic and Biodistribution of Baicalin-Loaded Liposome on Cerebral Ischemia-Reperfusion after iv. Administration in Rats. *Molecules*. 2018 Jul 17;23(7).
17. Luo YP, Zhang H, Hu HF, Cao ZY, Zhang XZ, Cao L, Wang ZZ, Xiao W. [Protective effects of Ginkgo Terpene Lactones Meglumine Injection on focal cerebral ischemia in rats]. *Aging Dis*. 2017 Dec 1;8(6):850-867.